

# TEBBIE FUNGICIDE

FOR SUPPRESSION OF FUSARIUM HEAD BLIGHT AND CONTROL OF  
MULTIPLE LEAF DISEASE IN NUMEROUS CROPS

**Crops:** Wheat, Barley, Oats, and Soybeans

**Diseases:** Fusarium Head Blight (S), Asian soybean rust, Crown rust, Frogeye leaf spot, Leaf rust, Stripe rust, Tan spot, Net blotch, Spot blotch, Powdery mildew, Scald, Leaf blotch, Glume blotch, & Stem rust

## Application Rates and Timing

Full Rate for Fusarium Head Blight — 200mL/ac

For optimal suppression: Apply Tebbie when 75% of wheat heads on main stem are fully emerged to when 50% of heads on main stem are in flower

3/4 Rate for Leaf Disease Control — 150 mL/ac

For optimal control: Apply Tebbie between the late vegetative stage to the end of the flowering stage

Active Ingredient  
Tebuconazole –Group 3

Formulation  
Suspension  
250 g/L

Package Size  
2 x 9.46L

Registration No. 33901

## Vandaele Ag Inputs

Bob Vandaele  
204-522-5613

Warren Alexander  
204-576-0118

Hayley Macaulay  
204-665-2384

